



## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:	PETRONIS	*	
Serial No.	10/516,406	*	
Filed:	December 1, 2004	*	Art Unit: 1617
For:	DETECTION OF EPIGENETIC ABNORMALITIES AND DIAGNOSTIC METHOD BASED THEREON	* * *	Examiner: (to be assigned)

## INFORMATION DISCLOSURE STATEMENT

To the Commissioner for Patents  
P. O. Box 1450  
Alexandria, VA 22313-1450

Dear Sir:

In accordance with 37 C.F.R. §1.56, applicant wishes to call the attention of the Examiner to the following references:

<u>Inventor(s)</u>	<u>U.S. Patent Nos.</u>
Mullis et al	4,683,195
Mullis	4,683,202
Mullis et al	4,800,159
Duffy	5,871,917

Foreign Patent Documents

<u>Country</u>	<u>Document No.</u>
PCT	US98/13818

Other Documents

Alitalo T, Tiihonen J, Hakola P, de la Chapelle A (1988): Molecular characterization of a Y;15 translocation segregating in a family. *Hum Genet* 79:29-35.

Altschul et al., *J. Mol. Biol.* (1990) 215:403-10

Bassett AS, Chow EW, Waterworth DM, Brzustowicz L (2001): Genetic insights into schizophrenia. *Can J. Psychiatry* 46:131-7.

Bennett RL, Karayiorgou M, Sobin CA, Norwood TH, Kay MA (1997): Identification of an interstitial deletion in an adult female with schizophrenia, mental retardation, and dysmorphic features: further support for a putative schizophrenia-susceptibility locus at 5q21-23.1. *Am J Hum Genet* 61:1450-4.

Berrettini W (2002): Review of bipolar molecular linkage and association studies. *Curr Psychiatry Rep* 4:124-9.

Berrettini WH 2000a): Are schizophrenic and bipolar disorders related? A review of family and molecular studies. *Biol Psychiatry* 48:531-8.

Berrettini WH 2000b): Susceptibility loci for bipolar disorder; overlap with inherited vulnerability to schizophrenia. *Biol Psychiatry* 47:245-51.

Bray NJ, Owen MJ. Searching for schizophrenia genes. *Trends Mol Med.* 2001; 7(4):169-74.

Brzustowicz LM, Hodgkinson KA, Chow EW, Honer WG, Bassett AS. Location of a major susceptibility locus for familial schizophrenia on chromosome 1q21-q22. *Science* 2000 Apr 28;288(5466):678-82.

Camp NJ, Neuhausen SL, Tiobech J, Polloi A, Coon H, Myles-Worsley M (2001): Genomewide multipoint linkage analysis of seven extended Palauan pedigrees with schizophrenia, by a Markov-chain Monte Carlo method. *Am J Hum Genet* 69:1278-89.

Cardno AG, Rijdsdijk FV, Sham PC, Murray RM, McGuffin P (2002): A twin study of genetic relationships between psychotic symptoms. *Am J Psychiatry* 159:539-45.

Chow EW, Bassett AS, Weksberg R. Velo-cardio-facial syndrome and psychotic disorders: implications for psychiatric genetics. *Am J Med Genet* 1994;54(2):107-12

DeLisi LE, Shaw SH, Crow TJ, et al (2002): A genome-wide scan for linkage to chromosomal regions in 382 sibling pairs with schizophrenia or schizoaffective disorder. *Am J Psychiatry* 159:803-12.

Detera-Wadleigh SD, Badner JA, Berrettini WH, et al (1999): A high-density genome scan detects evidence for a bipolar-disorder susceptibility locus on 13q32 and other potential loci on 1q32 and 18p11.2. *Proc Natl Acad Sci U S A* 96:5604-9

Detera-Wadleigh SC, Badner JA, Goldin LR, et al (1996): Affected-sib-pair analyses reveal support of prior evidence for a susceptibility locus for bipolar disorder, on 21q. *Am J Hum Genet* 58:1279-85.

Egeland et al. (1987) *Nature* 325: 783-787.

Evans KL, Brown J, Shibasaki Y, et al (1995): A contiguous clone map over 3 Mb on the long arm of chromosome 11 across a balanced translocation associated with schizophrenia. *Genomics* 28:420-8.

Ewald H, Wang AG, Vang M, Mors O, Nyegaard M, Kruse TA (1999): A haplotype-based study of lithium responding patients with bipolar affective disorder on the Faroe Islands. *Psychiatr Genet* 9:23-34.

Fan et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6223-27.

Faraone SV, Matise T, Svrakic D, et al (1998): Genome scan of European-American schizophrenia pedigrees: results of the NIMH Genetics Initiative and Millennium Consortium. *Am J Med Genet* 81:290-5.

Feinberg, A. P. and Vogelstein, B. (1983) Hypomethylation of ras oncogenes in primary human cancers. *Biochem Biophys Res Commun* 111, 47-54.

Feinberg, A. P., Gehrke, C. W., Kuo, K. C., and Ehrlich, M. (1988). Reduced genomic 5-methylcytosine content in human colonic neoplasia. *Cancer Res* 48, 1159-61.

Filippova GN, Thienes CP, Penn BH, et al (2001): CTCF-binding sites flank CTG/CAG repeats and form a methylation-sensitive insulator at the DM1 locus. *Nat Genet* 28:335-43.

Florl AR, Franke KH, Niederacher D, Gerharz CD, Seifert HH, Schulz WA. DNA methylation and the mechanisms of CDKN2A inactivation in transitional cell carcinoma of the urinary bladder. *Laboratory Investigation*, 80(10):1513-22, 2000.

Freimer NB, Reus VI, Escamilla MA, et al (1996): Genetic mapping using haplotype, association and linkage methods suggests a locus for severe bipolar disorder (BPI) at 18q22-q23. *Nat Genet* 12:436-41.

Gama-Sosa, M.A., Slagel, V. A., Trewyn, R. W., Oxenhandler, R., Kuo, K. C., Gehrke, C. W., and Ehrlich, M. (1983). The 5-methylcytosine content of DNA from human tumors. *Nucleic Acids Res* 11, 6883-94.

Gershon ES. Bipolar illness and schizophrenia as oligogenic diseases: implications for the future. *Biol Psychiatry*. 2000 Feb 1; 47(3):240-4.

Gill M, Vallada H, Collier D, et al (1996): A combined analysis of D22S278 marker alleles in affected sib-pairs: support for a susceptibility locus for schizophrenia at chromosome 22q12. Schizophrenia Collaborative Linkage Group (Chromosome 22). *Am J Med Genet* 67:40-5.

Gonzalzo, M.L. and Jones, P.A. (1997) Mutagenic and epigenetic effects of DNA methylation. *Mutat. Res.* 386(2), 107-18.

H. Suzuki et al., 6 *Nature Genet* 332-33 (1994) (please note that the specification incorrectly lists this reference as H. Suzuki et al., 7 *Nature Genet* 432-38 (1994))

Hedenfalk I, Duggan D, Chen Y, et al (2001): Gene-expression profiles in hereditary breast cancer. *N Engl J Med* 344:539-48.

Henikoff S, Matzke MA Exploring and explaining epigenetic effects. *Trends Genet* 1997; 13(8):293-5.

Hernandez I, Sokolov BP. Abnormalities in 5-HT<sub>2A</sub> receptor mRNA expression in frontal cortex of chronic elderly schizophrenics with varying histories of neuroleptic treatment. *J Neurosci Res*. 2000; 59(2):218-25.

Issa J. P., et al. (1994) *Nature Genetics* 7:536-40.

Johnston-Wilson NL, Sims CD, Hofmann JP, et al (2000): Disease-specific alterations in frontal cortex brain proteins in schizophrenia, bipolar disorder, and major depressive disorder. The Stanley Neuropathology Consortium. *Mol Psychiatry* 5:242-9.

Joo EJ, Lee JH, Cannon TD, Price RA (1999): Possible association between schizophrenia and a CAG repeat polymorphism in the spinocerebellar ataxia type 1 (SCA1) gene on human chromosome 6p23. *Psychiatr Genet* 9:7-11.

Jurgens B, Schmitz-Drager BJ, Schulz WA. Hypomethylation of L1 LINE sequences prevailing in human urothelial carcinoma. *Cancer Research*, 56(24):5698-703, 1996.

Karlsson H, Bachmann S, Schroder J, McArthur J, Torrey EF, Yolken RH (2001): Retroviral RNA identified in the cerebrospinal fluids and brains of individuals with schizophrenia. *Proc Natl Acad Sci U S A* 98:4634-9.

Kelsoe JR, Spence MA, Loetscher E, et al (2001): A genome survey indicates a possible susceptibility locus for bipolar disorder on chromosome 22. *Proc Natl Acad Sci U S A* 98:585-90.

Kendler KS, Myers JM, O'Neill FA, et al (2000): Clinical features of schizophrenia and linkage to chromosomes 5q, 6p, 8p, and 10p in the Irish Study of High-Density Schizophrenia Families. *Am J Psychiatry* 157:402-8.

Kwoh et al., 8 *Am. Biotechnol. Lab.* 14 (1990).

Laird et al., DNA Methylation and Cancer, 3 Human Molecular Genetics 1487, 1488 (1994).

Lapeyre, J. N. and Becker, F. F. (1979). 5-Methylcytosine content of nuclear DNA during chemical hepatocarcinogenesis and in carcinomas which result. *Biochem Biophys Res Commun* 87, 698-705.

Le Corre S, Harper CG, Lopez P, Ward P, Catts S (2000): Increased levels of expression of an NMDARI splice variant in the superior temporal gyrus in schizophrenia. *Neuroreport* 11:983-6.

Liang G, et al. (2002 Identification of DNA methylation differences during tumorigenesis by methylation-sensitive arbitrarily primed polymerase chain reaction. *Methods* 27(2): 150-5.

McNeil TF (1995): Perinatal risk factors and schizophrenia: selective review and methodological concerns. *Epidemiol Rev* 17:107-12.

Miniou P, Bourc'his D, Molina Gomes D, Jeanpierre M, Viegas-Pequignot E (1997): Undermethylation of Alu sequences in ICF syndrome: molecular and in situ analysis. *Cytogenet Cell Genet* 77:308-13.

Mirnics K, Middleton FA, Lewis DA, Levitt P (2001): Analysis of complex brain disorders with gene expression microarrays: schizophrenia as a disease of the synapse. *Trends Neurosci* 24:479-86.

Moises HW, Yang L, Li T, et al (1995): Potential linkage disequilibrium between schizophrenia and locus D22S278 on the long arm of chromosome 22. *Am J Med Genet* 60:465-7.

Mors O, Mortensen PB, Ewald H (2001): No evidence of increased risk for schizophrenia or bipolar affective disorder in persons with aneuploidies of the sex chromosomes. *Psychol Med* 31:425-30.

Mowry BJ, Nancarrow DJ (2001): Molecular genetics of schizophrenia. *Clin Exp Pharmacol Physiol* 28:66-9.

Docket No.: 04340-PCT-PA

Inventor: PETRONIS

Serial No.: 10/516,406

Mujaheed M, Corbex M, Lichtenberg P, et al (2000): Evidence for linkage by transmission disequilibrium test analysis of a chromosome 22 microsatellite marker D22S278 and bipolar disorder in a Palestinian Arab population. *Am J Med Genet* 96:836-8.

Myles-Worsley M, Coon H, McDowell J, et al (1999): Linkage of a composite inhibitory phenotype to a chromosome 22q locus in eight Utah families. *Am J Med Genet* 88:544-50.

Nurnberger JI Jr, Foroud T. Genetics of bipolar affective disorder. *Curr Psychiatry Rep* 2000 Apr;2(2):147-57.

Ogawa, et al., 362 *Nature* 749-51 (1993).

P. V. Pedone et al., 3 *Hum. Mol. Genet.* 1117-21 (1994).

Petit J, Boisseau P, Taine L, Gauthier B, Arveiler B (1999): A YAC contig encompassing the 11q14.3 breakpoint of a translocation associated with schizophrenia, and including the tyrosinase gene. *Mamm Genome* 10:649-52.

Petronis A, Gottesman, II, Crow TJ, et al (2000): Psychiatric epigenetics: a new focus for the new century. *Mol Psychiatry* 5:342-6.

Petronis A, Paterson AD, Kennedy JL. Schizophrenia: an epigenetic puzzle? *Schizophrenia Bulletin* 25:4: 639-655, 1999; Petronis A. The genes for major psychosis: aberrant sequence or regulation? *Neuropsychopharmacology*, 23(1):1-12; 2000.

Petronis A. Human morbid genetics revisited: relevance of epigenetics. *Trends Genet.* 2001 Mar;17(3):142-6.

Petronis A. The genes for major psychosis: aberrant sequence or regulation? *Neuropsychopharmacology*, 23(1):1-12; 2000.

Pfeifer GP, et al. (1989) *Science* 246:810-13.

Docket No.: 04340-PCT-PA

Inventor: PETRONIS

Serial No.: 10/516,406

Popendikyte V, Laurinavicius A, Paterson AD, Macciardi F, Kennedy JL, Petronis A. DNA methylation at the putative promotor region of the human dopamine D2 receptor gene. *Neuroreport* 1999; 10:1249-55.

Pulver AE, Karayiorgou M, Wolyniec PS, et al (1994): Sequential Strategy to identify a susceptibility gene for schizophrenia: report of potential linkage on chromosome 22q12-q13.1:Part 1. *Am J Med Genet* 54:36-43.

S. Rainier et al, 55 *Cancer Res.* 1836-38 (1995).

Rainier et al. 362 *Nature* 747-49 (1993).

Rakyan VK, Blewitt ME, Druker R, Preis JJ, Whitelaw E (2002): Metastable epialleles in mammals. *Trends Genet* 18:348-51.

Rice JP, Goate A, Williams JT, et al (1997): Initial genome scan of the NIMH genetics initiative bipolar pedigrees: chromosomes 1, 6, 8, 10, and 12. *Am J Med Genet* 74:247-53.

Robertson KD and Wolffe AP (2000) DNA methylation in health and disease. *Nature Review Genet* 1:11-9.

S. Zhan et al., 94 *J. Clin. Invest.* 445-48 (1994).

Saiki et al. (1988 *Science* 239:487-491).

Saluz and Jost (1989) *Proc Natl Acad Sci USA* 86:2602-6.

Schizophrenia Collaborative Linkage Group (1998): A transmission disequilibrium and linkage analysis of D22S278 marker alleles in 574 families: further support for a susceptibility locus for schizophrenia at 22q12. *Schizophr Res* 32:115-21.

Schwab SG, Albus M, Hallmayer J, et al (1995a): Evaluation of a susceptibility gene for schizophrenia on chromosome 6p by multipoint affected sib-pair linkage analysis. *Nat Genet* 11:325-7.



Schwab SG, Hallmayer J, Albus M, et al (1998): Further evidence for a susceptibility locus on chromosome 10p14-p11 in 72 families with schizophrenia by nonparametric linkage analysis. *Am J Med Genet* 81:302-7.

Schwab SG, Lerer B, Albus M, et al (1995b): Potential linkage for schizophrenia on chromosome 22q12-q13: a replication study. *Am J Med Genet* 60:436-43.

Sherrington R, Brynjolfsson J, Petursson H, Potter M, Dudleston K, Barraclough B, Wasmuth J, Dobbs M, Gurling H. Localization of a susceptibility locus for schizophrenia on chromosome 5. *Nature*. 1988 Nov 10;336(6195):164-7.

Sinnott D, Richer C, Deragon JM, Labuda D. Alu RNA transcripts in human embryonal carcinoma cells. Model of post-transcriptional selection of master sequences. *Journal of Molecular Biology*, 226(3):689-706, 1992.

St Clair D, Blackwood D, Muir W, Carothers A, Walker M, Spowart G, Gosden C, Evans HJ. Association within a family of a balanced autosomal translocation with major mental illness. *Lancet*. 1990 Jul 7;336(8706):13-6.

Straub RE, MacLean CJ, Martin RB, et al (1998): A schizophrenia locus may be located in region 10p15-p11. *Am J Med Genet* 81:296-301.

Straub RE, MacLean CJ, O'Neill FA, Walsh D, Kendler KS (1997): Support for a possible schizophrenia vulnerability locus in region 5q22-31 in Irish families. *Mol Psychiatry* 2:148-55.

Susser E, Neugebauer R, Hoek HW, et al (1996): Schizophrenia after prenatal famine. Further evidence. *Arch Gen Psychiatry* 53:25-31.

Thaker GK, Carpenter WT Jr. Advances in schizophrenia. *Nat Med*. 2001 Jun;7(6):667-71.

Tsuang MT, Stone WS, Faraone SV. Genes, environment and schizophrenia. *Br J Psychiatry Suppl*. 2001 Apr; 40:s18-24.

Docket No.: 04340-PCT-PA  
Inventor: PETRONIS  
Serial No.: 10/516,406

Verheyen GR, Villafuerte SM, Del-Favero J, et al (1999): Genetic refinement and physical mapping of a chromosome 18q candidate region for bipolar disorder. Eur J Hum Genet 7:427-34.

Wang S, Detera-Wadleigh SD, Coon H, et al (1996): Evidence of linkage disequilibrium between schizophrenia and the SCA1 CAG repeat on chromosome 6p23. Am J Hum Genet 59:731-6.

Xu G-L, Bestor TH. Nature Genetics 17:376-379, 1997.

Applicant respectfully submits that the above-mentioned references taken alone or in combination neither anticipate nor render obvious the present invention. Consideration of the foregoing and the making of the references formally of record in the instant application is respectfully requested.

Form PTO/SB/42 (6-95) is submitted for the convenience of the Examiner.

If there are any fees required, please charge our Deposit Account No. 02-2839.

Respectfully submitted,

Mar 30, 2006  
Date

Robert M. Gamson  
Robert M. Gamson  
Reg. No. 32,986  
Attorney for Applicant

RMG/jjr (03/30/06)  
Enclosures

ARMSTRONG, KRATZ, QUINTOS, HANSON & BROOKS, LLP  
502 Washington Avenue, Suite 220  
Towson, Maryland 21204  
Telephone: (410) 337-2295  
Fax: (410) 337-2296

P:\HOME\Clients\0148\0041\04340 IDS Petronis-Gowling.wpd

CERTIFICATE OF TRANSMISSION  
I hereby certify that this correspondence is being deposited with the U.S. Postal Service as Express Mail in an envelope addressed to: Commissioner for Patents, P. O. Box 1450, Alexandria, VA 22313-1450  
Date: March 30, 2006  
Express Label No.: EV 826565275 US  
By: Judy Robertson  
Judy Robertson

37 CFR 1.501(b)

## INFORMATION DISCLOSURE CITATION IN A PATENT

*(Use several sheets if necessary)*

**Docket Number: 04340-PCT-PA (0148.0041)**

Serial Number: 10/516,406

**Applicant: PETRONIS**

**Issue Date:**

Group Art Unit: 1617

## U.S. PATENT DOCUMENTS

[illegible]

	Document Number	Date	Country	Class	Subclass	Translation	
						Yes	No
	US98/13818	07/02/1998	PCT				

## OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)

	87 references . Please see attachment.

**EXAMINER**

Date Considered \_\_\_\_\_

**Burden Hour Statement:** This form is estimated to take 2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. **DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**

Other Documents

Alitalo T, Tiihonen J, Hakola P, de la Chapelle A (1988): Molecular characterization of a Y;15 translocation segregating in a family. Hum Genet 79:29-35.

Altschul et al., J. Mol. Biol. (1990) 215:403-10

Bassett AS, Chow EW, Waterworth DM, Brzustowicz L (2001): Genetic insights into schizophrenia. Can J. Psychiatry 46:131-7.

Bennett RL, Karayiorgou M, Sobin CA, Norwood TH, Kay MA (1997): Identification of an interstitial deletion in an adult female with schizophrenia, mental retardation, and dysmorphic features: further support for a putative schizophrenia-susceptibility locus at 5q21-23.1. Am J Hum Genet 61:1450-4.

Berrettini W (2002): Review of bipolar molecular linkage and association studies. Curr Psychiatry Rep 4:124-9.

Berrettini WH 2000a): Are schizophrenic and bipolar disorders related? A review of family and molecular studies. Biol Psychiatry 48:531-8.

Berrettini WH 2000b): Susceptibility loci for bipolar disorder; overlap with inherited vulnerability to schizophrenia. Biol Psychiatry 47:245-51.

Bray NJ, Owen MJ. Searching for schizophrenia genes. Trends Mol Med. 2001; 7(4):169-74.

Brzustowicz LM, Hodgkinson KA, Chow EW, Honer WG, Bassett AS. Location of a major susceptibility locus for familial schizophrenia on chromosome 1q21-q22. Science 2000 Apr 28;288(5466):678-82.

Camp NJ, Neuhausen SL, Tiobech J, Polloi A, Coon H, Myles-Worsley M (2001): Genomewide multipoint linkage analysis of seven extended Palauan pedigrees with schizophrenia, by a Markov-chain Monte Carlo method. Am J Hum Genet 69:1278-89.

Cardno AG, Rijsdijk FV, Sham PC, Murray RM, McGuffin P (2002): A twin study of genetic relationships between psychotic symptoms. Am J Psychiatry 159:539-45.

Chow EW, Bassett AS, Weksberg R. Velo-cardio-facial syndrome and psychotic disorders: implications for psychiatric genetics. *Am J Med Genet* 1994;54(2):107-12

DeLisi LE, Shaw SH, Crow TJ, et al (2002): A genome-wide scan for linkage to chromosomal regions in 382 sibling pairs with schizophrenia or schizoaffective disorder. *Am J Psychiatry* 159:803-12.

Detera-Wadleigh SD, Badner JA, Berrettini WH, et al (1999): A high-density genome scan detects evidence for a bipolar-disorder susceptibility locus on 13q32 and other potential loci on 1q32 and 18p11.2. *Proc Natl Acad Sci U S A* 96:5604-9

Detera-Wadleigh SC, Badner JA, Goldin LR, et al (1996): Affected-sib-pair analyses reveal support of prior evidence for a susceptibility locus for bipolar disorder, on 21q. *Am J Hum Genet* 58:1279-85.

Egeland et al. (1987) *Nature* 325: 783-787.

Evans KL, Brown J, Shibasaki Y, et al (1995): A contiguous clone map over 3 Mb on the long arm of chromosome 11 across a balanced translocation associated with schizophrenia. *Genomics* 28:420-8.

Ewald H, Wang AG, Vang M, Mors O, Nyegaard M, Kruse TA (1999): A haplotype-based study of lithium responding patients with bipolar affective disorder on the Faroe Islands. *Psychiatr Genet* 9:23-34.

Fan et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6223-27.

Faraone SV, Matisse T, Svrakic D, et al (1998): Genome scan of European-American schizophrenia pedigrees: results of the NIMH Genetics Initiative and Millennium Consortium. *Am J Med Genet* 81:290-5.

Feinberg, A. P. and Vogelstein, B. (1983) Hypomethylation of ras oncogenes in primary human cancers. *Biochem Biophys Res Commun* 111, 47-54.

Feinberg, A. P., Gehrke, C. W., Kuo, K. C., and Ehrlich, M. (1988). Reduced genomic 5-methylcytosine content in human colonic neoplasia. *Cancer Res* 48, 1159-61.

Filippova GN, Thienes CP, Penn BH, et al (2001): CTCF-binding sites flank CTG/CAG repeats and form a methylation-sensitive insulator at the DM1 locus. *Nat Genet* 28:335-43.

Florl AR, Franke KH, Niederacher D, Gerharz CD, Seifert HH, Schulz WA. DNA methylation and the mechanisms of CDKN2A inactivation in transitional cell carcinoma of the urinary bladder. *Laboratory Investigation*, 80(10):1513-22, 2000.

Freimer NB, Reus VI, Escamilla MA, et al (1996): Genetic mapping using haplotype, association and linkage methods suggests a locus for severe bipolar disorder (BPI) at 18q22-q23. *Nat Genet* 12:436-41.

Gama-Sosa, M.A., Slagel, V. A., Trewyn, R. W., Oxenhandler, R., Kuo, K. C., Gehrke, C. W., and Ehrlich, M. (1983). The 5-methylcytosine content of DNA from human tumors. *Nucleic Acids Res* 11, 6883-94.

Gershon ES. Bipolar illness and schizophrenia as oligogenic diseases: implications for the future. *Biol Psychiatry*. 2000 Feb 1; 47(3):240-4.

Gill M, Vallada H, Collier D, et al (1996): A combined analysis of D22S278 marker alleles in affected sib-pairs: support for a susceptibility locus for schizophrenia at chromosome 22q12. Schizophrenia Collaborative Linkage Group (Chromosome 22). *Am J Med Genet* 67:40-5.

Gonzalgo, M.L. and Jones, P.A. (1997) Mutagenic and epigenetic effects of DNA methylation. *Mutat. Res.* 386(2), 107-18.

H. Suzuki et al., 6 *Nature Genet* 332-33 (1994) (please note that the specification incorrectly lists this reference as H. Suzuki et al., 7 *Nature Genet* 432-38 (1994))

Hedenfalk I, Duggan D, Chen Y, et al (2001): Gene-expression profiles in hereditary breast cancer. *N Engl J Med* 344:539-48.

Henikoff S, Matzke MA Exploring and explaining epigenetic effects. *Trends Genet* 1997; 13(8):293-5.

Hernandez I, Sokolov BP. Abnormalities in 5-HT<sub>2A</sub> receptor mRNA expression in frontal cortex of chronic elderly schizophrenics with varying histories of neuroleptic treatment. *J Neurosci Res.* 2000; 59(2):218-25.

Issa J. P., et al. (1994) *Nature Genetics* 7:536-40.

Johnston-Wilson NL, Sims CD, Hofmann JP, et al (2000): Disease-specific alterations in frontal cortex brain proteins in schizophrenia, bipolar disorder, and major depressive disorder. The Stanley Neuropathology Consortium. *Mol Psychiatry* 5:242-9.

Joo EJ, Lee JH, Cannon TD, Price RA (1999): Possible association between schizophrenia and a CAG repeat polymorphism in the spinocerebellar ataxia type 1 (SCA1) gene on human chromosome 6p23. *Psychiatr Genet* 9:7-11.

Jurgens B, Schmitz-Drager BJ, Schulz WA. Hypomethylation of L1 LINE sequences prevailing in human urothelial carcinoma. *Cancer Research*, 56(24):5698-703, 1996.

Karlsson H, Bachmann S, Schroder J, McArthur J, Torrey EF, Yolken RH (2001): Retroviral RNA identified in the cerebrospinal fluids and brains of individuals with schizophrenia. *Proc Natl Acad Sci U S A* 98:4634-9.

Kelsoe JR, Spence MA, Loetscher E, et al (2001): A genome survey indicates a possible susceptibility locus for bipolar disorder on chromosome 22. *Proc Natl Acad Sci U S A* 98:585-90.

Kendler KS, Myers JM, O'Neill FA, et al (2000): Clinical features of schizophrenia and linkage to chromosomes 5q, 6p, 8p, and 10p in the Irish Study of High-Density Schizophrenia Families. *Am J Psychiatry* 157:402-8.

Kwoh et al., 8 *Am. Biotechnol. Lab.* 14 (1990).

Laird et al., *DNA Methylation and Cancer*, 3 *Human Molecular Genetics* 1487, 1488 (1994).

Lapeyre, J. N. and Becker, F. F. (1979). 5-Methylcytosine content of nuclear DNA during chemical hepatocarcinogenesis and in carcinomas which result. *Biochem Biophys Res Commun* 87, 698-705.

Le Corre S, Harper CG, Lopez P, Ward P, Catts S (2000): Increased levels of expression of an NMDAR1 splice variant in the superior temporal gyrus in schizophrenia. *Neuroreport* 11:983-6.

Liang G, et al. (2002 Identification of DNA methylation differences during tumorigenesis by methylation-sensitive arbitrarily primed polymerase chain reaction. *Methods* 27(2): 150-5.

McNeil TF (1995): Perinatal risk factors and schizophrenia: selective review and methodological concerns. *Epidemiol Rev* 17:107-12.

Miniou P, Bourc'his D, Molina Gomes D, Jeanpierre M, Viegas-Pequignot E (1997): Undermethylation of Alu sequences in ICF syndrome: molecular and in situ analysis. *Cytogenet Cell Genet* 77:308-13.

Mirnics K, Middleton FA, Lewis DA, Levitt P (2001): Analysis of complex brain disorders with gene expression microarrays: schizophrenia as a disease of the synapse. *Trends Neurosci* 24:479-86.

Moises HW, Yang L, Li T, et al (1995): Potential linkage disequilibrium between schizophrenia and locus D22S278 on the long arm of chromosome 22. *Am J Med Genet* 60:465-7.

Mors O, Mortensen PB, Ewald H (2001): No evidence of increased risk for schizophrenia or bipolar affective disorder in persons with aneuploidies of the sex chromosomes. *Psychol Med* 31:425-30.

Mowry BJ, Nancarrow DJ (2001): Molecular genetics of schizophrenia. *Clin Exp Pharmacol Physiol* 28:66-9.

Mujaheed M, Corbex M, Lichtenberg P, et al (2000): Evidence for linkage by transmission disequilibrium test analysis of a chromosome 22 microsatellite marker D22S278 and bipolar disorder in a Palestinian Arab population. *Am J Med Genet* 96:836-8.

Myles-Worsley M, Coon H, McDowell J, et al (1999): Linkage of a composite inhibitory phenotype to a chromosome 22q locus in eight Utah families. *Am J Med Genet* 88:544-50.

Nurnberger JI Jr, Foroud T. Genetics of bipolar affective disorder. *Curr Psychiatry Rep* 2000 Apr;2(2):147-57.

Ogawa, et al., 362 *Nature* 749-51 (1993).

P. V. Pedone et al., 3 *Hum. Mol. Genet.* 1117-21 (1994).

Petit J, Boisseau P, Taine L, Gauthier B, Arveiler B (1999): A YAC contig encompassing the 11q14.3 breakpoint of a translocation associated with schizophrenia, and including the tyrosinase gene. *Mamm Genome* 10:649-52.

Petronis A, Gottesman, II, Crow TJ, et al (2000): Psychiatric epigenetics: a new focus for the new century. *Mol Psychiatry* 5:342-6.

Petronis A, Paterson AD, Kennedy JL. Schizophrenia: an epigenetic puzzle? *Schizophrenia Bulletin* 25:4: 639-655, 1999; Petronis A. The genes for major psychosis: aberrant sequence or regulation? *Neuropsychopharmacology*, 23(1):1-12; 2000.



Petronis A. Human morbid genetics revisited: relevance of epigenetics. Trends Genet. 2001 Mar;17(3):142-6.

Petronis A. The genes for major psychosis: aberrant sequence or regulation? Neuropsychopharmacology, 23(1):1-12; 2000.

Pfeifer GP, et al. (1989) Science 246:810-13.

Popendikyte V, Laurinavicius A, Paterson AD, Macciardi F, Kennedy JL, Petronis A. DNA methylation at the putative promotor region of the human dopamine D2 receptor gene. Neuroreport 1999; 10:1249-55.

Pulver AE, Karayiorgou M, Wolyniec PS, et al (1994): Sequential Strategy to identify a susceptibility gene for schizophrenia: report of potential linkage on chromosome 22q12-q13.1:Part 1. Am J Med Genet 54:36-43.

S. Rainier et al, 55 Cancer Res. 1836-38 (1995).

Rainner et al. 362 Nature 747-49 (1993).

Rakyan VK, Blewitt ME, Druker R, Preis JJ, Whitelaw E (2002): Metastable epialleles in mammals. Trends Genet 18:348-51.

Rice JP, Goate A, Williams JT, et al (1997): Initial genome scan of the NIMH genetics initiative bipolar pedigrees: chromosomes 1, 6, 8, 10, and 12. Am J Med Genet 74:247-53.

Robertson KD and Wolffe AP (2000) DNA methylation in health and disease. Nature Review Genet 1:11-9.

S. Zhan et al., 94 J. Clin. Invest. 445-48 (1994).

Saiki et al. (1988 Science 239:487-491).

Saluz and Jost (1989) Proc Natl Acad Sci USA 86:2602-6.

Schizophrenia Collaborative Linkage Group (1998): A transmission disequilibrium and linkage analysis of D22S278 marker alleles in 574 families: further support for a susceptibility locus for schizophrenia at 22q12. Schizophr Res 32:115-21.

Schwab SG, Albus M, Hallmayer J, et al (1995a): Evaluation of a susceptibility gene for schizophrenia on chromosome 6p by multipoint affected sib-pair linkage analysis. *Nat Genet* 11:325-7.

Schwab SG, Hallmayer J, Albus M, et al (1998): Further evidence for a susceptibility locus on chromosome 10p14-p11 in 72 families with schizophrenia by nonparametric linkage analysis. *Am J Med Genet* 81:302-7.

Schwab SG, Lerer B, Albus M, et al (1995b): Potential linkage for schizophrenia on chromosome 22q12-q13: a replication study. *Am J Med Genet* 60:436-43.

Sherrington R, Brynjolfsson J, Petursson H, Potter M, Dudleston K, Barracough B, Wasmuth J, Dobbs M, Gurling H. Localization of a susceptibility locus for schizophrenia on chromosome 5. *Nature*. 1988 Nov 10;336(6195):164-7.

Sinnett D, Richer C, Deragon JM, Labuda D. Alu RNA transcripts in human embryonal carcinoma cells. Model of post-transcriptional selection of master sequences. *Journal of Molecular Biology*, 226(3):689-706, 1992.

St Clair D, Blackwood D, Muir W, Carothers A, Walker M, Spowart G, Gosden C, Evans HJ. Association within a family of a balanced autosomal translocation with major mental illness. *Lancet*. 1990 Jul 7;336(8706):13-6.

Straub RE, MacLean CJ, Martin RB, et al (1998): A schizophrenia locus may be located in region 10p15-p11. *Am J Med Genet* 81:296-301.

Straub RE, MacLean CJ, O'Neill FA, Walsh D, Kendler KS (1997): Support for a possible schizophrenia vulnerability locus in region 5q22-31 in Irish families. *Mol Psychiatry* 2:148-55.

Susser E, Neugebauer R, Hoek HW, et al (1996): Schizophrenia after prenatal famine. Further evidence. *Arch Gen Psychiatry* 53:25-31.

Thaker GK, Carpenter WT Jr. Advances in schizophrenia. *Nat Med*. 2001 Jun;7(6):667-71.

Tsuang MT, Stone WS, Faraone SV. Genes, environment and schizophrenia. *Br J Psychiatry Supl*. 2001 Apr; 40:s18-24.

Verheyen GR, Villafuerte SM, Del-Favero J, et al (1999): Genetic refinement and physical mapping of a chromosome 18q candidate region for bipolar disorder. *Eur J Hum Genet* 7:427-34.

Wang S, Detera-Wadleigh SD, Coon H, et al (1996): Evidence of linkage disequilibrium between schizophrenia and the SCA1 CAG repeat on chromosome 6p23. Am J Hum Genet 59:731-6.

Xu G-L, Bestor TH. Nature Genetics 17:376-379, 1997.